

Discussion #5

- 1. Starburst/AGN diagnostics: detection**
- 2. Starburst/AGN diagnostics: quantitative**
- 3. Impact of AGN on starburst and vice-versa**

Starburst/AGN Diagnostics: Detection


- **How reliable are OIR diagnostics?**
 - **In dusty systems?**
(what are the many optical composites among ULIRGs?)

Starburst/AGN Diagnostics: Detection

- **How reliable are OIR diagnostics?**
 - **In dusty systems?**
(what are the many optical composites among ULIRGs?)
- **What about MIR?**
 - **What is the cause of the (often) low (MIR line/FIR) ratios in many ULIRGs and some IR-bright QSOs?**
 - **Density, radiation field, covering factor?**

Method

- 6 diagnostics: $[O\ IV]/[Ne\ II]$, $[Ne\ V]/[Ne\ II]$, $EW(PAH)$, $L(MIR)/L(FIR)$, $15/5\ \mu m$ ratio, $30/15\ \mu m$ ratio...
- All agree to within $\pm 10\text{-}15\%$ on average



Results


- Strong correlations with optical spectral types!
- AGN% increases along merger sequence, but substantial stochastic variations

(SV, Rupke, et al. 2009a)

Starburst/AGN Diagnostics: Detection

- **How reliable are OIR diagnostics?**
 - In dusty systems?
(what are the many optical composites among ULIRGs?)
- **What about MIR?**
 - What is the cause of the low (line/FIR) ratios?
 - Density, radiation field, covering factor?
- **Can FIR-Submm-mm help?**
 - CO ladder → PDR vs XDR

CO Ladder: PDR vs XDR




(van der Werf et al. 2010)

Starburst/AGN Diagnostics: Detection

- **How reliable are OIR diagnostics?**
 - In dusty systems?
(what are the many optical composites among ULIRGs?)
- **What about MIR?**
 - What is the cause of the low (line/FIR) ratios?
 - Density, radiation field, covering factor?
- **Can FIR-Submm-mm help?**
 - CO ladder → PDR vs XDR
- **How useful is the radio (if there's no jet)?**
 - Minor bolometric contributor, free-free absorption,
 - Is monitoring essential?
- **How useful are the hard X-rays?**
 - Sensitive to absorption/scattering along line of sight to nucleus

Chandra/XMM/Suzaku Results




- $\langle \text{AGN}\% \rangle \sim L(2-10 \text{ keV}) / L_{\text{BOL}}$
- Generally good agreement with Spitzer results:
 - Strong correlations with optical spectral types
 - AGN% increases along merger sequence, but with substantial stochastic variations


(Teng & SV 2010, submitted)

Starburst/AGN Diagnostics: Quantitative

- **L[O III] as an indicator of $dM/dt(BH)$?**
 - **SB contamination...**
 - **Dust extinction**
- **MIR: [O IV], [Ne V]...**
- **How useful are the hard X-rays?**
 - **Absorption/scattering correction**
 - **Bolometric correction**
- **How useful is the radio? (SB vs radio-quiet AGN)**
 - **Large bolometric correction**
 - **Free-free absorption**



(Wild et al. 2010; LaMassa et al. 2010)



(Winter et al. 2010)

AGN – Starburst Connection

- **AGN – SB connection or AGN – SF connection?**
- **Selection effects: bimodal distributions of SFH**
- **Use of fundamental plane / what do latest results imply?**
- **What are the observational/theoretical prospects of understanding inner ~ 100 pc?**
 - **Bar-within-bar, nuclear spirals, stochastic process?**
 - **Feeding the AGN with 0.1% of SB material, AGB winds?**
 - **Stellar population = $f(L_{\text{AGN}})$???**
 - **AGN = $f(\text{SB})$???**